# 2track separation study with GARFIELD++

Yasuyuki Akiba, Itaru Nakagawa, Gaku Mitsuka, Yorito Yamaguchi July 29<sup>th</sup>, 2016


## Setup (HV & Gas)

♦ HV configuration

Anode: 0V, Cathode: -4500V, Field: -2600V, Guard: -800V,

Channel: -1500V


- $\Leftrightarrow$  Gas: Ar-C<sub>2</sub>H<sub>6</sub> (50:50)
  - ✓ Gas properties are calculated by Magboltz
- ♦ X: mirror periodicity, Y: GND at upper & lower boundaries
- ♦ Lorentz angle: OFF (on-going)



## Signal from anode wire


- ♦ Induced current, i(t) is calculated using Shockley-Ramo theorem
  - $\checkmark i(t) = -q\mathbf{v} \cdot \mathbf{E}_{w}(\mathbf{r})$ , q: charge, v: drift velocity
- ♦ Convolution with response function of shaping amp
  - $\checkmark$  V(t)=(t/n $\tau$ )<sup>n</sup>e<sup>n-t/ $\tau$ </sup>


t=2ns, n=3 from PHENIX DC electronics



### 2track simulation

- ♦ Input tracks
  - ✓ Signal: electron ( $p_T$ =5GeV/c,  $\alpha$ =2deg)
  - ✓ BG:  $\pi$  (p<sub>T</sub>=0.5GeV/c,  $\alpha$ = +/-20deg)
  - ✓ Only drift electrons generated by ionization, no multiplication
- $\Leftrightarrow$  Electron track always passes at  $(x_e, y_e) = (1, 0.3)$ , and p track at  $(x_\pi, y_\pi) = (1 \Delta x, 0.3)$ .
  - $\checkmark$  Note: s-6 at  $(x_{s-6}, y_{s-6}) = (0,0.3)$  [cm]
  - ✓ Variation of  $\Delta x$  :  $\Delta x$  = 0.08, 0.10, 0.12, 0.14, 0.16 cm






# Peak finding

- ♦ Procedure of peak finding
  - 1. Peak finding using TSpectrum class
  - 2. Gaussian fit with peak positions from TSpectrum
    - a. If npeaks==2, double Gaussian fit
    - b. If npeaks==1, single Gaussian fit
      - i. If peak width of fit > 7ns, double Gaussian fit

#### Averaged signal shapes of 2tracks at s-6





# Examples of peak finding


♦ Examples of succeeded & failed events

✓ BG:  $\pi$ ,  $p_T$ =500MeV,  $\Delta x$ =0.10cm

✓ Track trajectory :










$$\Delta(x_e-x)$$

- $\Leftrightarrow$  BG:  $\pi$ ,  $p_T$ =500MeV
- $\Rightarrow$  Distance between e &  $\pi$ :  $\Delta x = 0.08, 0.10, 0.12, 0.14, 0.16 cm$
- $\diamond$  Width of fitted Gaussian for e peak:  $\sigma_e$ <0.1mm



## Summary table (BG: $\pi$ , p<sub>T</sub>=500MeV)






	2peaks detected by Tspectrum	1peak detected by Tspectrum	peak width>7ns	peak width≦7ns	x <sub>e</sub> -x <0.2mm	x <sub>e</sub> -x <0.2mm &&  x <sub>pi</sub> -x <0.3mm
∆x=0.08cm	2	98	94	4	73(2)	60
	4	96	93	3	83(2)	69
∆x=0.10cm	18	82	77	5	76(3)	60
	14	86	81	5	72(4)	54
∆x=0.12cm	42	58	50	8	88(8)	59
	21	79	68	11	83(10)	54
∆x=0.14cm	55	45	35	10	93(10)	59
	50	50	38	12	87(10)	54
∆x=0.16cm	69	31	22	9	93(6)	63
	69	31	23	8	95(8)	65

## Detection probability of electron peak

- $\Rightarrow$  Definition of detection probability of electron peak : events with  $\Delta(x_e-x)<0.2$ mm/100 events
- $\diamondsuit$  Probability > 0.7 ( $\Delta$ x>0.08cm with 0.5GeV/c  $\pi$  track)



## To-do

- ♦ Simulation with Lorentz angle ON is now ongoing.
  - ✓ Need more time for finding good wire configuration

